

Losartan Suspension 2.5 mg/mL

INGREDIENTS:

Losartan 50 mg tablet	6 tablets
Purified water USP	6 mL
Ora-Plus/Ora-Sweet SF*	QSAD: 120 mL

EQUIPMENT AND SUPPLIES:

Graduated cylinder

PREPARATION DETAILS:

1. Place tablets in an appropriately sized amber bottle.
2. Add purified water to amber bottle and shake bottle for 2 minutes.
3. Let solution stand for 1 hour and then shake for 1 minute to disperse the tablet contents.
4. Add 114 mL of vehicle and shake well for 1 minute to disperse contents.

Special Instructions — *Mix 60 mL of Ora-Plus with 60 mL of Ora-Sweet SF. Use mixture as vehicle or use Ora-Blend SF.

Quality-Control Procedures — Visually inspect for physical appearance of formulation and container closure integrity (no leakage, cracks in container, or improper seals).

Labeling Requirements — Extemporaneously compounded preparation. For oral use only. Store in refrigerator. Shake well before use.

Storage Conditions/Stability — Refrigerate. Stable for 28 days.

STABILITY STUDY DETAILS:

Study Container Type — Amber polyethylene terephthalate (PET) bottle

Referenced Manufacturers — Losartan tablets (Cozaar, Merck & Co); Ora-Plus, Ora-Sweet SF (Paddock Laboratories, LLC).

Stability-Indicating Study — No

REFERENCE

1. Cozaar [prescribing information]. Whitehouse Station, NJ: Merck & Co; 2019.