

FUTURE TREATMENTS

The future of treatment options for dyslipidemia will most likely involve the evaluation of other parameters besides LDL-C. Advanced lipid testing that evaluates LDL particle size, HDL particle size, direct measured LDL, homocysteine, lipoprotein (a), and CRP are gaining popularity. Additionally, the measurement of apoB may eventually overtake LDL as the primary monitoring parameter for dyslipidemia. ApoB is the primary lipoprotein in LDL-C and is indicative of the amount of small dense LDL particles that are known to be more atherogenic and are more commonly present in patients with CHD, than that of the large buoyant LDL particles. Some of the new innovative drug therapies such as torcetrapib (cholesteryl ester transfer protein inhibitor) and lapaquistat (a squalene synthase inhibitor) have failed to reach the market due to increases in mortality and liver toxicity, respectively. However, another cholesteryl ester transfer protein inhibitor, anacetrapib is still being investigated. Darapladib, a selective lipoprotein-associated phospholipase A2 (Lp-PLA2) inhibitor, is also being studied. Lp-PLA2 is a marker of vascular inflammation, produced by macrophages, within the atherosclerotic plaque itself. This substance attaches to LDL-C and is responsible for the hydrolysis of oxidized phospholipids on LDL-C particles. It is considered an emerging risk factor for CHD.

The most promising drugs to lower LDL levels are inhibitors of proprotein convertase subtilisin/kexin type 9 (PCSK9). Normally, LDL receptors bind LDL and internalize it for subsequent lysosomal degradation, and then the receptors return to the plasma membrane to repeat the process. The enzyme PCSK9 prevents LDL receptors from returning to the cell surface by facilitating their degradation. Monoclonal antibodies (evolocumab, alirocumab, and bococizumab) that inhibit PCSK9 activity are in various stages of new drug development. Successful inhibition of PCSK9 allows continuous LDL receptor recycling and dramatic reductions in plasma LDL levels.

REFERENCES

1. Expert panel on detection, evaluation, and treatment of high blood cholesterol in adults. Executive summary of the third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III). *JAMA*. 2001;285:2486-97.
2. Lloyd-Jones D, Adams R, Carnethon M, et al. Heart disease and stroke statistics 2009 update: a report from the American Heart Association Statistics Committee and Stroke Subcommittee. *Circulation*. 2009;119:e21-181.
3. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 1—executive summary. *J Clin Lipidol*. 2014;8:473-488.
4. Expert panel on detection, evaluation, and treatment of high blood cholesterol in adults. Executive summary of the third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (adult treatment panel III). *Arch Intern Med*. 1988;148:36-69.
5. Grundy SM, Cleeman JI, Merz CNB, et al. NCEP report: implications of recent clinical trials for the national cholesterol education program adult treatment panel III guidelines. *Circulation*. 2004;110:227-39.
6. Stone NJ, Robinson JG, Lichtenstein AH. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association task force on practice guidelines. *Circulation*. 2014;129(25 Suppl 2):S1-S45. Epub 2013 Nov 12.
7. Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. *Circulation*. 2002;105:1135-43.
8. Cayette AJ, Palacino JJ, Horten K, et al. Chronic inhibition of nitric oxide production accelerates neointima formation and impairs endothelial function in hypercholesterolemic rabbits. *Arterioscler Thromb*. 1994;14:753-59.
9. Steinberg D, Parthasarathy S, Carew TE, et al. Beyond cholesterol. Modifications of low-density lipoprotein that increase its atherogenicity. *N Engl J Med*. 1989;320:915-24.
10. Nathan CF. Secretory products of macrophages. *J Clin Invest*. 1987;79:319-26.

11. Cockerill GW, Rye KA, Gamble JR, et al. High-density lipoproteins inhibit cytokine-induced expression of endothelial cell adhesion molecules. *Arterioscler Thromb Vasc Biol.* 1995;15:1987-94.
12. D'Agostino RB Sr. Understanding the cardiovascular disease risk functions—aim, development, and evaluation. *N Engl J Med.* 2014; 370(17):1652-53 (inset).
13. Goff DC Jr, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association task force on practice guidelines. *Circulation.* 2013 November 12.
14. Standards of medical care in diabetes-2014. *Diabetes Care.* 2014;37 (Supplement 1):S14-S80.
15. Mantel-Teeuwisse AK, Kloosterman JM, Maitland-van der Zee A, et al. Drug-induced lipid changes: a review of the unintended effects of some commonly used drugs on serum lipid levels. *Drug Safety.* 2001;24:443-46.
16. Atmaca M, Kuloglu M, Tezcan E, et al. Serum leptin and triglyceride levels in patients on treatment with atypical antipsychotics. *J Clin Psychiatry.* 2003;64:598-604.
17. Moore R, Hernandez D, Valantine H. Calcineurin inhibitors and post-transplant hyperlipidemias. *Drug Safety.* 2001;24:755-66.
18. Talbert R. Hyperlipidemia. In *Pharmacotherapy: A Pathophysiologic Approach, 5th ed.* New York, NY: McGraw-Hill; 2002:395-417.
19. McKenney JM. Pharmacotherapy of dyslipidemia. *Cardiovasc Drugs Ther.* 2001;15:413-22.
20. Merck/Schering-Plough Pharmaceuticals. Ezetimibe (Zetia) product insert. North Wales, PA; May 2009.
21. Daiichi Sankyo, Inc. Colesevelam hydrochloride (Welchol) product insert. Parsippany, NJ; January 2008.
22. Duval C, Chinetti G, Trottein F, et al. The role of PPARs in atherosclerosis. *Trends Mol Med.* 2002;8:422-30.
23. Hottelart C, El Elper N, Rose F, et al. Fenofibrate increases creatinemia by increasing metabolic production of creatinine. *Nephron.* 2002;92:536-41.
24. Levin A, Duncan L, Djurdjev O, et al. A randomized placebo-controlled double blind trial of lipid lowering strategies in patients with renal insufficiency: diet modification with or without fenofibrate. *Clin Nephrol.* 2000;53:140-46.
25. Deighan CJ, Caslake MJ, McConnell M, et al. Comparative effects of cerivastatin and fenofibrate on the atherogenic lipoprotein phenotype in proteinuric renal disease. *J Am Soc Nephrol.* 2001;12:341-48.
26. Dalton RN, Crimet D, Ansquer JC. The effect of fenofibrate on glomerular filtration rate (GFR) and other renal function tests: a double-blind placebo-controlled cross-over study in healthy subjects. Philadelphia: American Society of Nephrology 2005.
27. Hottelart C, el Esper N, Achard JM, et al. Fenofibrate increases blood creatinine, but does not change the glomerular filtration rate in patients with mild renal insufficiency. *Nephron.* 1999;20:41-4.
28. Abbott Laboratories. Niacin (Niaspan) product insert. North Chicago, IL; April 2009.
29. Merck & Co., Inc. Lovastatin (Mevacor) product insert. Whitehouse Station, NJ; September 2008.
30. Reliant Pharmaceuticals, LLC. Omega-3-acid ethyl esters 90 (Lovaza) product insert. Liberty Corner, NJ; June 2008.
31. Rojas-Fernandez CH, Goldstein LB, Levey AI, et al. An assessment by the Statin Cognitive Safety Task Force: 2014 update. *J Clin Lipidol.* 2014;8:S5-S16.
32. Maki KC, Ridker PM, Brown WV, et al. An assessment by the Statin Diabetes Safety Task Force: 2014 update. *J Clin Lipidol.* 2014;8:S17-S29.
33. Gotto AM. Safety and statin therapy: reconsidering the risks and benefits. *Arch Intern Med.* 2003;163:657-59.
34. Cohen DE, Anania FA, Chalasani N. An assessment of stain safety by hepatologists. *Am J Cardiol.* 2006;97:77C-81C.
35. Bays H, Cohen DE, Chalasani N, et al. An assessment by the Statin Liver Safety Task Force: 2014 update. *J Clin Lipidol.* 2014;8:S47-S57.
36. Law M, Rudnicka AR. Statin safety: a systematic review. *Am J Cardiol.* 2006;97:S52-S60.
37. Merck & Co., Inc. Simvastatin (Zocor) product insert. Whitehouse Station, NJ; June 2008.
38. AstraZeneca Pharmaceuticals LP. Rosuvastatin (Crestor) product insert. Wilmington, DE; February 2009.
39. Alsheikh-Ali AA, Duvin JT, Karas RH. Risk of adverse events with fibrates. *Am J Cardiol.* 2004;94:935-38.
40. Rosenson RS, Baker SK, Jacobson TA, et al. An assessment by the Statin Muscle Safety Task Force: 2014 update. *J Clin Lipidol.* 2014;8:S58-S71.